

WAC 16-324-398 Field inspection disease tolerance. (1) Compliance with a 0.0% tolerance is not intended, nor should it be construed, to mean that the lot inspected is free from the disease. It means that the disease was not detected during visual inspections of the seed lot.

(2) First and second field inspection tolerances, expressed as percentages.

Factor	Nuclear	G 1	G 2	G 3	G 4	G 5
Varietal mixture	0.10	0.10	0.20	0.30	0.40	0.50
Mosaic	0.00	0.10	0.25	0.50	0.75	1.00
Leafroll	0.00	0.10	0.25	0.30	0.50	0.50
Total visible virus	0.00	0.20	0.50	0.80	1.25	1.50
Phytoplasmas (including <i>Candidatus liberibacter</i>)	0.00	0.00	0.10	0.20	0.50	1.00
Black leg	0.00	0.10	0.50	1.00	2.00	*
Ring rot	0.00	0.00	0.00	0.00	0.00	0.00
Nematode	0.00	0.00	0.00	0.00	0.00	0.00
Spindle tuber viroid and other quarantined pests	0.00	0.00	0.00	0.00	0.00	0.00

*Tolerance for black leg does not apply to G5.

[Statutory Authority: RCW 15.14.015 and chapter 34.05 RCW. WSR 15-01-042, § 16-324-398, filed 12/9/14, effective 1/9/15; WSR 13-12-014, § 16-324-398, filed 5/24/13, effective 6/24/13. Statutory Authority: Chapters 15.13, 15.14 and 34.05 RCW. WSR 04-12-026, § 16-324-398, filed 5/26/04, effective 6/26/04. Statutory Authority: Chapter 15.14 RCW. WSR 02-12-010, § 16-324-398, filed 5/23/02, effective 6/23/02; WSR 00-20-070, § 16-324-398, filed 10/3/00, effective 11/3/00. Statutory Authority: RCW 15.14.030. WSR 97-11-028, § 16-324-398, filed 5/14/97, effective 6/14/97.]